

# Quartic and Octic Characters Modulo $n$

STEVEN FINCH

December 21, 2009

**ABSTRACT.** The average number of primitive quadratic Dirichlet characters of modulus  $n$  tends to a constant as  $n \rightarrow \infty$ . The same is true for primitive cubic characters. It is therefore surprising that, as  $n \rightarrow \infty$ , the average number of primitive quartic characters of modulus  $n$  grows with  $\ln(n)$ , and that the average number of primitive octic characters of modulus  $n$  grows with  $\ln(n)^2$ . Leading coefficients in the asymptotic expressions are also computed.

Let  $\mathbb{Z}_n^*$  denote the group (under multiplication modulo  $n$ ) of integers relatively prime to  $n$ , and let  $\mathbb{C}^*$  denote the group (under ordinary multiplication) of nonzero complex numbers. We wish to count homomorphisms  $\chi : \mathbb{Z}_n^* \rightarrow \mathbb{C}^*$  satisfying certain requirements. A Dirichlet character  $\chi$  is quadratic if  $\chi(k)^2 = 1$  for every  $k$  in  $\mathbb{Z}_n^*$ . It is well-known that

$$\begin{array}{l} \# \text{ quadratic Dirichlet characters} \\ \text{of modulus } \leq N \end{array} = \sum_{n \leq N} a(n) \sim \frac{6}{\pi^2} N \ln(N)$$

as  $N \rightarrow \infty$ , where  $a(n)$  is multiplicative with

$$a(2^r) = \begin{cases} 1 & \text{if } r = 1, \\ 2 & \text{if } r = 2, \\ 4 & \text{if } r \geq 3, \end{cases} \quad a(p^r) = 2$$

for prime  $p \geq 3$  and  $r \geq 1$ ; and

$$\begin{array}{l} \# \text{ primitive quadratic Dirichlet} \\ \text{characters of modulus } \leq N \end{array} = \sum_{n \leq N} b(n) \sim \frac{6}{\pi^2} N$$

where  $b(n)$  is multiplicative with

$$b(2^r) = \begin{cases} 0 & \text{if } r = 1, \\ 1 & \text{if } r = 2, \\ 2 & \text{if } r = 3, \\ 0 & \text{if } r \geq 4, \end{cases} \quad b(p^r) = \begin{cases} 1 & \text{if } r = 1, \\ 0 & \text{if } r \geq 2 \end{cases}$$

---

<sup>0</sup> Copyright © 2009 by Steven R. Finch. All rights reserved.

for prime  $p \geq 3$ . Is it surprising that the same constant  $6/\pi^2$  appears in connection with averages of both  $a(n)$  and  $b(n)$ ? No. The underlying structure linking these two sequences is the Möbius inversion formula [1]:

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^s} = \zeta(s) \sum_{n=1}^{\infty} \frac{b(n)}{n^s} \quad \text{for } \operatorname{Re}(s) > c \text{ for some } c < 1.$$

This can be proved directly here. On the one hand [2],

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{a(n)}{n^s} &= \left(1 + \frac{1}{2^s} + \frac{2}{2^{2s}} + \sum_{r=3}^{\infty} \frac{4}{2^{rs}}\right) \prod_{p>2} \left(1 + \sum_{r=1}^{\infty} \frac{2}{p^{rs}}\right) \\ &= \left(1 + \frac{1}{2^s} + \frac{2}{2^{2s}} + \frac{4}{2^s - 1} - \frac{4}{2^s} - \frac{4}{2^{2s}}\right) \prod_{p>2} \left(1 + \frac{2}{p^s - 1}\right) \\ &= \left(1 - \frac{3}{2^s} - \frac{2}{2^{2s}} + \frac{4}{2^s - 1}\right) \prod_{p>2} \left(1 - \frac{1}{p^s}\right)^{-2} \left(1 - \frac{1}{p^{2s}}\right) \\ &= \left(1 - \frac{3}{2^s} - \frac{2}{2^{2s}} + \frac{4}{2^s - 1}\right) \left(1 - \frac{1}{2^s}\right)^2 \zeta(s)^2 \left(1 - \frac{1}{2^{2s}}\right)^{-1} \zeta(2s)^{-1} \\ &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}}\right) \left(1 - \frac{1}{2^s}\right) \zeta(s)^2 \left(1 - \frac{1}{2^{2s}}\right)^{-1} \zeta(2s)^{-1}; \end{aligned}$$

on the other hand,

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{b(n)}{n^s} &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}}\right) \prod_{p>2} \left(1 + \frac{1}{p^s}\right) \\ &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}}\right) \prod_{p>2} \left(1 - \frac{1}{p^s}\right)^{-1} \left(1 - \frac{1}{p^{2s}}\right) \\ &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}}\right) \left(1 - \frac{1}{2^s}\right) \zeta(s) \left(1 - \frac{1}{2^{2s}}\right)^{-1} \zeta(2s)^{-1}. \end{aligned}$$

The leading coefficient  $6/\pi^2$  arises because

$$\left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}}\right) \left(1 - \frac{1}{2^s}\right) \left(1 - \frac{1}{2^{2s}}\right)^{-1} \zeta(2s)^{-1} \Big|_{s=1} = \frac{1}{\zeta(2)} = \frac{6}{\pi^2}.$$

The Möbius inversion formula is valid, in fact, for arbitrary  $\ell^{\text{th}}$ -order Dirichlet characters. From the preceding case  $\ell = 2$ , we observe the evaluation of  $\sum_{n=1}^{\infty} b(n)n^{-s}$  to be slightly less complicated than that for  $\sum_{n=1}^{\infty} a(n)n^{-s}$ . Hence, to simplify calculations, our focus will be on  $b(n)$  for the cases  $\ell = 3, 4, 8$ .

## 1. CUBIC CHARACTERS

**1.1. General.** A Dirichlet character  $\chi$  is cubic if  $\chi(k)^3 = 1$  for every  $k$  in  $\mathbb{Z}_n^*$ . We have [3]

$$a(2^r) = 1, \quad a(3^r) = \begin{cases} 1 & \text{if } r = 1, \\ 3 & \text{if } r \geq 2, \end{cases} \quad a(p^r) = \begin{cases} 1 & \text{if } p \equiv 2 \pmod{3}, \\ 3 & \text{if } p \equiv 1 \pmod{3} \end{cases}$$

for prime  $p \geq 5$  and  $r \geq 1$ . The asymptotics for  $\sum_{n \leq N} a(n)$ , as well as the coefficient, were studied in [4] (via a different proof than the following).

**1.2. Primitive.** We have

$$b(2^r) = 0, \quad b(3^r) = \begin{cases} 2 & \text{if } r = 2, \\ 0 & \text{otherwise,} \end{cases} \quad b(p^r) = \begin{cases} 2 & \text{if } r = 1 \text{ \& } p \equiv 1 \pmod{3}, \\ 0 & \text{otherwise} \end{cases}$$

for prime  $p \geq 5$  and  $r \geq 1$ . The asymptotics for  $\sum_{n \leq N} b(n)$ , as well as the coefficient, were studied in [5]. We obtain

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{b(n)}{n^s} &= \left(1 + \frac{2}{3^{2s}}\right) \prod_{p \equiv 1 \pmod{3}} \left(1 + \frac{2}{p^s}\right) \\ &= \left(1 + \frac{2}{3^{2s}}\right) \prod_{p \equiv 1 \pmod{3}} \left(1 - \frac{1}{p^s}\right)^{-2} \left(1 - \frac{1}{p^{2s}}\right) \left(1 - \frac{2}{p^s(p^s + 1)}\right) \end{aligned}$$

and it is known from section [4.1] that

$$\prod_{p \equiv 1 \pmod{3}} \left(1 - \frac{1}{p^s}\right)^{-2} \sim \zeta(s) \frac{\sqrt{3}}{2\pi} \prod_{p \equiv 1 \pmod{3}} \left(1 - \frac{1}{p^2}\right)^{-1} \quad (1)$$

as  $s \rightarrow 1$ ; after cancellation, the coefficient becomes

$$\frac{11}{9} \frac{\sqrt{3}}{2\pi} \prod_{p \equiv 1 \pmod{3}} \left(1 - \frac{2}{p(p+1)}\right) = 0.3170565167922841205670156....$$

Just as in the case of primitive quadratic characters, the function  $\zeta(s)$  appears with exponent 1. Hence by the Selberg-Delange method [4], the average number of primitive cubic characters of modulus  $n$  tends to a constant 0.317... as  $n \rightarrow \infty$ .

## 2. QUARTIC CHARACTERS

**2.1. General.** A Dirichlet character  $\chi$  is quartic (biquadratic) if  $\chi(k)^4 = 1$  for every  $k$  in  $\mathbb{Z}_n^*$ . We have [6]

$$a(2^r) = \begin{cases} 1 & \text{if } r = 1, \\ 2 & \text{if } r = 2, \\ 4 & \text{if } r = 3, \\ 8 & \text{if } r \geq 4, \end{cases} \quad a(p^r) = \begin{cases} 2 & \text{if } p \equiv 3 \pmod{4}, \\ 4 & \text{if } p \equiv 1 \pmod{4} \end{cases}$$

for prime  $p \geq 3$  and  $r \geq 1$ .

**2.2. Primitive.** We have

$$b(2^r) = \begin{cases} 1 & \text{if } r = 2, \\ 2 & \text{if } r = 3, \\ 4 & \text{if } r = 4, \\ 0 & \text{otherwise,} \end{cases} \quad b(p^r) = \begin{cases} 1 & \text{if } r = 1 \text{ \& } p \equiv 3 \pmod{4}, \\ 3 & \text{if } r = 1 \text{ \& } p \equiv 1 \pmod{4} \\ 0 & \text{otherwise} \end{cases}$$

for prime  $p \geq 3$  and  $r \geq 1$ . We obtain

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{b(n)}{n^s} &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}} + \frac{4}{2^{4s}}\right) \prod_{p \equiv 3 \pmod{4}} \left(1 + \frac{1}{p^s}\right) \cdot \prod_{p \equiv 1 \pmod{4}} \left(1 + \frac{3}{p^s}\right) \\ &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}} + \frac{4}{2^{4s}}\right) \prod_{p \equiv 3 \pmod{4}} \left(1 - \frac{1}{p^s}\right)^{-1} \left(1 - \frac{1}{p^{2s}}\right) \\ &\quad \cdot \prod_{p \equiv 1 \pmod{4}} \left(1 - \frac{1}{p^s}\right)^{-3} \left(1 - \frac{1}{p^{2s}}\right) \left(1 - \frac{5p^s - 3}{p^{2s}(p^s + 1)}\right) \\ &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}} + \frac{4}{2^{4s}}\right) \left(1 - \frac{1}{2^s}\right) \zeta(s) \left(1 - \frac{1}{2^{2s}}\right)^{-1} \zeta(2s)^{-1} \\ &\quad \cdot \prod_{p \equiv 1 \pmod{4}} \left(1 - \frac{1}{p^s}\right)^{-2} \left(1 - \frac{5p^s - 3}{p^{2s}(p^s + 1)}\right) \end{aligned}$$

and, from section [4.2],

$$\prod_{p \equiv 1 \pmod{4}} \left(1 - \frac{1}{p^s}\right)^{-2} \sim \zeta(s) \frac{1}{\pi} \prod_{p \equiv 1 \pmod{4}} \left(1 - \frac{1}{p^2}\right)^{-1} \sim \zeta(s) \frac{\pi}{16K^2} \quad (2)$$

as  $s \rightarrow 1$ ; thus the coefficient becomes

$$\frac{7}{\pi} \frac{1}{16K^2} \prod_{p \equiv 1 \pmod{4}} \left(1 - \frac{5p - 3}{p^2(p + 1)}\right) = 0.1908767211685284480112237....$$

where  $K$  is the Landau-Ramanujan constant [7]. Unlike the preceding cases, the function  $\zeta(s)$  appears with exponent 2. Hence by the Selberg-Delange method [4], the average number of primitive quartic characters of modulus  $n$  is asymptotically  $(0.190...) \ln(n)$  as  $n \rightarrow \infty$ . Is it surprising that the quartic case differs so dramatically from both the quadratic and cubic cases? We believe yes. There is no *a priori* reason for quartic characters to outnumber quadratic/cubic characters in such a manner.

## 3. OCTIC CHARACTERS

**3.1. General.** A Dirichlet character  $\chi$  is octic if  $\chi(k)^8 = 1$  for every  $k$  in  $\mathbb{Z}_n^*$ . We have

$$a(2^r) = \begin{cases} 1 & \text{if } r = 1, \\ 2 & \text{if } r = 2, \\ 4 & \text{if } r = 3, \\ 8 & \text{if } r = 4, \\ 16 & \text{if } r \geq 5, \end{cases} \quad a(p^r) = \begin{cases} 2 & \text{if } p \equiv 3 \pmod{4}, \\ 4 & \text{if } p \equiv 5 \pmod{8}, \\ 8 & \text{if } p \equiv 1 \pmod{8} \end{cases}$$

for prime  $p \geq 3$  and  $r \geq 1$ .

**3.2. Primitive.** We have

$$b(2^r) = \begin{cases} 1 & \text{if } r = 2, \\ 2 & \text{if } r = 3, \\ 4 & \text{if } r = 4, \\ 8 & \text{if } r = 5, \\ 0 & \text{otherwise,} \end{cases} \quad b(p^r) = \begin{cases} 1 & \text{if } r = 1 \text{ \& } p \equiv 3 \pmod{4}, \\ 3 & \text{if } r = 1 \text{ \& } p \equiv 5 \pmod{8}, \\ 7 & \text{if } r = 1 \text{ \& } p \equiv 1 \pmod{8}, \\ 0 & \text{otherwise} \end{cases}$$

for prime  $p \geq 3$  and  $r \geq 1$ . We obtain

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{b(n)}{n^s} &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}} + \frac{4}{2^{4s}} + \frac{8}{2^{5s}}\right) \prod_{\substack{p \equiv 3 \\ \pmod{4}}} \left(1 + \frac{1}{p^s}\right) \cdot \prod_{\substack{p \equiv 5 \\ \pmod{8}}} \left(1 + \frac{3}{p^s}\right) \cdot \prod_{\substack{p \equiv 1 \\ \pmod{8}}} \left(1 + \frac{7}{p^s}\right) \\ &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}} + \frac{4}{2^{4s}} + \frac{8}{2^{5s}}\right) \prod_{\substack{p \equiv 3 \\ \pmod{4}}} \left(1 - \frac{1}{p^s}\right)^{-1} \left(1 - \frac{1}{p^{2s}}\right) \\ &\quad \cdot \prod_{\substack{p \equiv 5 \\ \pmod{8}}} \left(1 - \frac{1}{p^s}\right)^{-3} \left(1 - \frac{1}{p^{2s}}\right) \left(1 - \frac{5p^s - 3}{p^{2s}(p^s + 1)}\right) \\ &\quad \cdot \prod_{\substack{p \equiv 1 \\ \pmod{8}}} \left(1 - \frac{1}{p^s}\right)^{-7} \left(1 - \frac{1}{p^{2s}}\right) \left(1 - \frac{27p^{5s} - 85p^{4s} + 125p^{3s} - 99p^{2s} + 41p^s - 7}{p^{6s}(p^s + 1)}\right) \\ &= \left(1 + \frac{1}{2^{2s}} + \frac{2}{2^{3s}} + \frac{4}{2^{4s}} + \frac{8}{2^{5s}}\right) \left(1 - \frac{1}{2^s}\right) \zeta(s) \left(1 - \frac{1}{2^{2s}}\right)^{-1} \zeta(2s)^{-1} \\ &\quad \cdot \prod_{\substack{p \equiv 5 \\ \pmod{8}}} \left(1 - \frac{1}{p^s}\right)^{-2} \left(1 - \frac{5p^s - 3}{p^{2s}(p^s + 1)}\right) \\ &\quad \cdot \prod_{\substack{p \equiv 1 \\ \pmod{8}}} \left(1 - \frac{1}{p^s}\right)^{-6} \left(1 - \frac{27p^{5s} - 85p^{4s} + 125p^{3s} - 99p^{2s} + 41p^s - 7}{p^{6s}(p^s + 1)}\right) \end{aligned}$$

and, from section [4.3],

$$\prod_{p \equiv 5 \pmod{8}} \left(1 - \frac{1}{p^s}\right)^{-4} \sim \zeta(s) \frac{1}{2 \ln(1 + \sqrt{2})} \prod_{p \equiv 5 \pmod{8}} \left(1 - \frac{1}{p^2}\right)^{-2}, \quad (3)$$

$$\prod_{p \equiv 1 \pmod 8} \left(1 - \frac{1}{p^s}\right)^{-4} \sim \zeta(s) \frac{2 \ln(1 + \sqrt{2})}{\pi^2} \prod_{p \equiv 1 \pmod 8} \left(1 - \frac{1}{p^2}\right)^{-2} \quad (4)$$

as  $s \rightarrow 1$ . An expression for the coefficient becomes clear. More importantly, the function  $\zeta(s)$  appears with exponent  $1 + 1/2 + 3/2 = 3$ . Hence by the Selberg-Delange method [4], the average number of primitive octic characters of modulus  $n$  has growth rate  $\approx \ln(n)^2$  as  $n \rightarrow \infty$ .

#### 4. EULER PRODUCT RESIDUES

Formulas (1), (2), (4) await proof, while the truth of (3) depends on both (2) and (4). Our approach uses the seemingly-unrelated method of Shanks & Schmid [8] for computing various generalized Landau-Ramanujan constants  $\kappa_m$ .

Fix an integer  $m \neq 0$ . Define

$$L_d(s) = \sum_{n=1}^{\infty} \left(\frac{d}{n}\right) \frac{1}{n^s}$$

where  $d = -m$  if  $4 \mid m$  and  $d = -4m$  otherwise;  $(\cdot/\cdot)$  is the Kronecker-Jacobi-Legendre symbol. Define also

$$\Lambda_m(s) = \sum_{n=1}^{\infty} \frac{f(n)}{n^s}$$

where  $f(n) = 1$  if there exist integers  $x, y$  such that  $n = x^2 + m y^2$  and  $f(n) = 0$  otherwise. Let

$$\kappa_m = \delta_m \sqrt{\frac{L_d(1)}{\pi} \frac{2|m|}{\varphi(2|m|)}} \prod_{\substack{\text{odd } p: \\ (\frac{-m}{p}) = -1}} \left(1 - \frac{1}{p^2}\right)^{-1/2}$$

where the rational number  $\delta_m$  is unspecified for the moment, and  $\varphi$  denotes the Euler totient function. It turns out that  $\sum_{n \leq N} f(n) \sim \kappa_m N / \sqrt{\ln(N)}$  as  $N \rightarrow \infty$ , although this fact is not directly relevant to our purposes.

**4.1. Case when  $\ell = 3$ .** Let  $m = 3$  and  $\delta_m = 2/3$ . It follows that

$$\begin{aligned} \kappa_3 &= \frac{2}{3} \sqrt{L_{-12}(1) \frac{1}{\pi} \frac{6}{\varphi(6)}} \prod_{\substack{\text{odd } p: \\ p \equiv 2 \pmod 3}} \left(1 - \frac{1}{p^2}\right)^{-1/2} \\ &= \frac{2}{3} \sqrt{\frac{\pi}{2\sqrt{3}} \frac{1}{\pi} \frac{6}{2}} \prod_{p \equiv 2 \pmod 3} \left(1 - \frac{1}{p^2}\right)^{-1/2} \cdot \left(1 - \frac{1}{2^2}\right)^{1/2} \\ &= \frac{1}{\sqrt{2\sqrt{3}}} \prod_{p \equiv 2 \pmod 3} \left(1 - \frac{1}{p^2}\right)^{-1/2} = 2^{1/2} 3^{-7/4} \pi \prod_{p \equiv 1 \pmod 3} \left(1 - \frac{1}{p^2}\right)^{1/2}. \end{aligned}$$

Now,

$$\Lambda_3(s) = (1 - 3^{-s})^{-1} \prod_{\substack{p \equiv 1 \\ \text{mod } 3}} (1 - p^{-s})^{-1} \cdot \prod_{\substack{p \equiv 2 \\ \text{mod } 3}} (1 - p^{-2s})^{-1}$$

by definition and

$$\Lambda_3(s)^2 = \zeta(s) L_{-3}(s) (1 - 3^{-s})^{-1} \prod_{\substack{p \equiv 2 \\ \text{mod } 3}} (1 - p^{-2s})^{-1}$$

by elementary considerations [9]. Dividing the second by the first, we have

$$\Lambda_3(s) = \zeta(s) L_{-3}(s) \prod_{\substack{p \equiv 1 \\ \text{mod } 3}} (1 - p^{-s})$$

and thus

$$\underbrace{\Lambda_3(s)(s-1)^{1/2}}_{\downarrow \sqrt{\pi} \kappa_3} = \underbrace{\zeta(s)(s-1)}_{\downarrow 1} \underbrace{L_{-3}(s)}_{\downarrow \frac{\pi}{3\sqrt{3}}} \prod_{\substack{p \equiv 1 \\ \text{mod } 3}} (1 - p^{-s}) \cdot (s-1)^{-1/2}$$

as  $s \rightarrow 1$ . Hence formula (1) is true.

**4.2. Case when  $\ell = 4$ .** Let  $m = 1$  and  $\delta_m = 1$ . It follows that

$$\begin{aligned} \kappa_1 &= 1 \sqrt{L_{-4}(1) \frac{1}{\pi} \frac{2}{\varphi(2)}} \prod_{\substack{\text{odd } p: \\ p \equiv 3 \text{ mod } 4}} \left(1 - \frac{1}{p^2}\right)^{-1/2} \\ &= \sqrt{\frac{\pi}{4} \frac{1}{\pi} \frac{2}{1}} \prod_{p \equiv 3 \text{ mod } 4} \left(1 - \frac{1}{p^2}\right)^{-1/2} \\ &= \frac{1}{\sqrt{2}} \prod_{p \equiv 3 \text{ mod } 4} \left(1 - \frac{1}{p^2}\right)^{-1/2} = K = \frac{\pi}{4} \prod_{p \equiv 1 \text{ mod } 4} \left(1 - \frac{1}{p^2}\right)^{1/2}. \end{aligned}$$

Now,

$$\Lambda_1(s) = (1 - 2^{-s})^{-1} \prod_{\substack{p \equiv 1 \\ \text{mod } 4}} (1 - p^{-s})^{-1} \cdot \prod_{\substack{p \equiv 3 \\ \text{mod } 4}} (1 - p^{-2s})^{-1}$$

by definition and

$$\Lambda_1(s)^2 = \zeta(s) L_{-4}(s) (1 - 2^{-s})^{-1} \prod_{\substack{p \equiv 3 \\ \text{mod } 4}} (1 - p^{-2s})^{-1}$$

by elementary considerations [10]. Dividing the second by the first, we obtain

$$\underbrace{\Lambda_1(s)(s-1)^{1/2}}_{\downarrow \sqrt{\pi}\kappa_1} = \underbrace{\zeta(s)(s-1)}_{\downarrow 1} \underbrace{L_{-4}(s)}_{\downarrow \frac{\pi}{4}} \prod_{\substack{p \equiv 1 \\ \text{mod } 4}} (1-p^{-s}) \cdot (s-1)^{-1/2}$$

as  $s \rightarrow 1$ . Hence formula (2) is true.

**4.3. Case when  $\ell = 8$ .** The argument here seems fairly roundabout and we wonder if a simpler approach is possible. Two values of  $m$  require consideration here. First, let  $m = 2$  and  $\delta_m = 1$ . It follows that

$$\begin{aligned} \kappa_2 &= 1 \sqrt{L_{-8}(1) \frac{1}{\pi} \frac{4}{\varphi(4)}} \prod_{\substack{\text{odd } p: \\ p \equiv 5, 7 \text{ mod } 8}} \left(1 - \frac{1}{p^2}\right)^{-1/2} \\ &= \sqrt{\frac{\pi}{2\sqrt{2}} \frac{1}{\pi} \frac{4}{2}} \prod_{p \equiv 5, 7 \text{ mod } 8} \left(1 - \frac{1}{p^2}\right)^{-1/2} \\ &= \frac{1}{\sqrt[4]{2}} \prod_{p \equiv 5, 7 \text{ mod } 8} \left(1 - \frac{1}{p^2}\right)^{-1/2} = 2^{-7/4} \pi \prod_{p \equiv 1, 3 \text{ mod } 8} \left(1 - \frac{1}{p^2}\right)^{1/2}. \end{aligned}$$

As before,

$$\begin{aligned} \Lambda_2(s) &= (1-2^{-s})^{-1} \prod_{\substack{p \equiv 1, 3 \\ \text{mod } 8}} (1-p^{-s})^{-1} \cdot \prod_{\substack{p \equiv 5, 7 \\ \text{mod } 8}} (1-p^{-2s})^{-1}, \\ \Lambda_2(s)^2 &= \zeta(s) L_{-8}(s) (1-2^{-s})^{-1} \prod_{\substack{p \equiv 5, 7 \\ \text{mod } 8}} (1-p^{-2s})^{-1} \end{aligned}$$

and, upon division, we obtain

$$\underbrace{\Lambda_2(s)(s-1)^{1/2}}_{\downarrow \sqrt{\pi}\kappa_2} = \underbrace{\zeta(s)(s-1)}_{\downarrow 1} \underbrace{L_{-8}(s)}_{\downarrow \frac{\pi}{2\sqrt{2}}} \prod_{\substack{p \equiv 1, 3 \\ \text{mod } 8}} (1-p^{-s}) \cdot (s-1)^{-1/2}.$$



Second, let  $m = -2$  and  $\delta_m = 1$ . It follows that

$$\begin{aligned}
 \kappa_{-2} &= 1 \sqrt{L_8(1) \frac{1}{\pi} \frac{4}{\varphi(4)}} \prod_{\substack{\text{odd } p: \\ p \equiv 3, 5 \pmod{8}}} \left(1 - \frac{1}{p^2}\right)^{-1/2} \\
 &= \sqrt{\frac{\sqrt{2} \ln(1 + \sqrt{2})}{\pi}} \prod_{p \equiv 3, 5 \pmod{8}} \left(1 - \frac{1}{p^2}\right)^{-1/2} \\
 &= 2^{-5/4} \ln(1 + \sqrt{2})^{1/2} \pi^{1/2} \prod_{p \equiv 1, 7 \pmod{8}} \left(1 - \frac{1}{p^2}\right)^{1/2}.
 \end{aligned}$$

As before,

$$\begin{aligned}
 \Lambda_{-2}(s) &= (1 - 2^{-s})^{-1} \prod_{\substack{p \equiv 1, 7 \\ \pmod{8}}} (1 - p^{-s})^{-1} \cdot \prod_{\substack{p \equiv 3, 5 \\ \pmod{8}}} (1 - p^{-2s})^{-1}, \\
 \Lambda_{-2}(s)^2 &= \zeta(s) L_8(s) (1 - 2^{-s})^{-1} \prod_{\substack{p \equiv 3, 5 \\ \pmod{8}}} (1 - p^{-2s})^{-1}
 \end{aligned}$$

and, upon division, we obtain

$$\underbrace{\Lambda_{-2}(s)(s-1)^{1/2}}_{\downarrow \sqrt{\pi} \kappa_{-2}} = \underbrace{\zeta(s)(s-1)}_{\downarrow 1} \underbrace{L_8(s)}_{\downarrow \frac{\ln(1+\sqrt{2})}{\sqrt{2}}} \prod_{\substack{p \equiv 1, 7 \\ \pmod{8}}} (1 - p^{-s}) \cdot (s-1)^{-1/2}.$$

Therefore

$$\begin{aligned}
 \lim_{s \rightarrow 1} \prod_{\substack{p \equiv 1, 3 \\ \pmod{8}}} \left(1 - \frac{1}{p^s}\right)^{-2} \cdot (s-1) &= \frac{\pi}{8} \frac{1}{\kappa_2^2} = \frac{2^{1/2}}{\pi} \prod_{\substack{p \equiv 1, 3 \\ \pmod{8}}} \left(1 - \frac{1}{p^2}\right)^{-1}, \\
 \lim_{s \rightarrow 1} \prod_{\substack{p \equiv 1, 7 \\ \pmod{8}}} \left(1 - \frac{1}{p^s}\right)^{-2} \cdot (s-1) &= \frac{\ln(1 + \sqrt{2})^2}{2\pi} \frac{1}{\kappa_{-2}^2} = \frac{2^{3/2} \ln(1 + \sqrt{2})}{\pi^2} \prod_{\substack{p \equiv 1, 7 \\ \pmod{8}}} \left(1 - \frac{1}{p^2}\right)^{-1}
 \end{aligned}$$

but these alone do not go far enough. A slightly revised formula (2):

$$\lim_{s \rightarrow 1} \prod_{\substack{p \equiv 1, 5 \\ \pmod{8}}} \left(1 - \frac{1}{p^s}\right)^{-2} \cdot (s-1) = \frac{\pi}{16} \frac{1}{\kappa_1^2} = \frac{1}{\pi} \prod_{\substack{p \equiv 1, 5 \\ \pmod{8}}} \left(1 - \frac{1}{p^2}\right)^{-1},$$

when coupled with the preceding two limits, make possible the isolation of  $p \equiv 1 \pmod{8}$  as follows:

$$\begin{aligned}
& \frac{4}{\pi^4} \ln(1 + \sqrt{2}) \overbrace{\prod_{1,3,5,7} (1 - p^{-2})^{-1}}^{=\pi^2/8} \cdot \prod_1 (1 - p^{-2})^{-2} \\
&= \frac{2^{1/2}}{\pi} \prod_{1,3} (1 - p^{-2})^{-1} \cdot \frac{1}{\pi} \prod_{1,5} (1 - p^{-2})^{-1} \cdot \frac{2^{3/2} \ln(1 + \sqrt{2})}{\pi^2} \prod_{1,7} (1 - p^{-2})^{-1} \\
&= \lim_{s \rightarrow 1} \left( \prod_{1,3} (1 - p^{-s})^{-2} \cdot (s-1) \right) \left( \prod_{1,5} (1 - p^{-s})^{-2} \cdot (s-1) \right) \left( \prod_{1,7} (1 - p^{-s})^{-2} \cdot (s-1) \right) \\
&= \lim_{s \rightarrow 1} \underbrace{\left( \prod_{1,3,5,7} (1 - p^{-s})^{-2} \cdot (s-1)^2 \right)}_{=1/4} \cdot \lim_{s \rightarrow 1} \left( \prod_1 (1 - p^{-s})^{-4} \cdot (s-1) \right).
\end{aligned}$$

Hence formula (4) is true.

## 5. UNANSWERED QUESTIONS

For the cases  $\ell = 5, 6, 7$ , the function  $\zeta(s)$  appears with exponents 1, 3, 1 respectively and [8]

$$\begin{aligned}
\kappa_{-3} &= \frac{1}{2} \sqrt{\frac{\sqrt{3} \ln(2+\sqrt{3})}{\pi}} \prod_{\substack{p \equiv 5,7 \\ \pmod{12}}} \left(1 - \frac{1}{p^2}\right)^{-1/2}, \\
\kappa_5 &= \frac{1}{2} \sqrt{\frac{\sqrt{5}}{2}} \prod_{\substack{p \equiv 11,13,17,19 \\ \pmod{20}}} \left(1 - \frac{1}{p^2}\right)^{-1/2}, \quad \kappa_{-5} = \frac{1}{2} \sqrt{\frac{\sqrt{5} \ln(9+4\sqrt{5})}{3\pi}} \prod_{\substack{p \equiv 2,3 \\ \pmod{5}}} \left(1 - \frac{1}{p^2}\right)^{-1/2}, \\
\kappa_6 &= \frac{1}{2} \sqrt{\frac{\sqrt{6}}{2}} \prod_{\substack{p \equiv 13,17,19,23 \\ \pmod{24}}} \left(1 - \frac{1}{p^2}\right)^{-1/2}, \quad \kappa_{-6} = \frac{1}{2} \sqrt{\frac{\sqrt{6} \ln(5+2\sqrt{6})}{2\pi}} \prod_{\substack{p \equiv 7,11,13,17 \\ \pmod{24}}} \left(1 - \frac{1}{p^2}\right)^{-1/2}, \\
\kappa_7 &= \frac{3}{4} \sqrt{\frac{\sqrt{7}}{6}} \prod_{\substack{p \equiv 3,5,6 \\ \pmod{7}}} \left(1 - \frac{1}{p^2}\right)^{-1/2}, \quad \kappa_{-7} = \frac{1}{2} \sqrt{\frac{\sqrt{7} \ln(8+3\sqrt{7})}{3\pi}} \prod_{\substack{p \equiv 5,11,13, \\ 15,17,23 \\ \pmod{28}}} \left(1 - \frac{1}{p^2}\right)^{-1/2}.
\end{aligned}$$

It does not seem to be possible to isolate  $p \equiv 1 \pmod{\ell}$  beyond the following partial results:

$$\begin{aligned}
\lim_{s \rightarrow 1} \prod_{\substack{p \equiv 1,4 \\ \pmod{5}}} \left(1 - \frac{1}{p^s}\right)^{-2} \cdot (s-1) &= \frac{\ln(9+4\sqrt{5})^2}{45\pi} \frac{1}{\kappa_{-5}^2} = \frac{\sqrt{5} \ln(9+4\sqrt{5})}{3\pi^2} \prod_{\substack{p \equiv 1,4 \\ \pmod{5}}} \left(1 - \frac{1}{p^2}\right)^{-1}, \\
\lim_{s \rightarrow 1} \prod_{\substack{p \equiv 1,2,4 \\ \pmod{7}}} \left(1 - \frac{1}{p^s}\right)^{-2} \cdot (s-1) &= \frac{9\pi}{112\pi} \frac{1}{\kappa_7^2} = \frac{3\sqrt{7}}{4\pi} \prod_{\substack{p \equiv 1,2,4 \\ \pmod{7}}} \left(1 - \frac{1}{p^2}\right)^{-1}
\end{aligned}$$

for  $\ell = 5, 7$ , which are deduced from

$$\Lambda_{-5}(s) = (1 - 5^{-s})^{-1} \prod_{\substack{p \equiv 1, 4 \\ \pmod{5}}} (1 - p^{-s})^{-1} \cdot \prod_{\substack{p \equiv 2, 3 \\ \pmod{5}}} (1 - p^{-2s})^{-1},$$

$$\Lambda_{-5}(s)^2 = \zeta(s) L_{20}(s) (1 - 5^{-s})^{-1} \prod_{\substack{p \equiv 2, 3 \\ \pmod{5}}} (1 - p^{-2s})^{-1} \cdot (1 + 2^{-s})^{-1}$$

and

$$\Lambda_7(s) = (1 - 7^{-s})^{-1} \prod_{\substack{2 < p \equiv 1, 2, 4 \\ \pmod{7}}} (1 - p^{-s})^{-1} \cdot \prod_{\substack{p \equiv 3, 5, 6 \\ \pmod{7}}} (1 - p^{-2s})^{-1} \cdot \left[ (1 - 2^{-s})^{-1} - 2^{-s} \right],$$

$$\Lambda_7(s)^2 = \zeta(s) L_{-28}(s) (1 - 7^{-s})^{-1} \cdot \prod_{\substack{p \equiv 3, 5, 6 \\ \pmod{7}}} (1 - p^{-2s})^{-1} \cdot \left[ (1 - 2^{-s})^{-1} - 2^{-s} \right]^2 (1 - 2^{-s})$$

as before. The L-series  $\Lambda_{-3}(s), \Lambda_5(s), \Lambda_6(s), \Lambda_{-6}(s), \Lambda_{-7}(s)$  might be difficult to study, due to a failure of  $f(n)$ -multiplicativity. When  $m = 6$ , for example,  $f(10) = 1$  since  $10 = 2^2 + 6 \cdot 1^2$ , but  $f(2) = f(5) = 0$ . When  $m = -6$ , as another example,  $f(10) = 1$  since  $10 = 4^2 - 6 \cdot 1^2$ , but again  $f(2) = f(5) = 0$ . Although the asymptotics of  $\sum_{n \leq N} b(n)$  for  $\ell = 5, 6, 7$  are understood, expressions for leading coefficients (analogous to those for  $\ell = 3, 4, 8$ ) remain open.

## 6. ACKNOWLEDGEMENT

I thank Pascal Sebah, my coauthor in [4], for his skillful numerical computations over many years! There is a way to avoid the “roundabout” calculation of residues (involving [8], as presented here) and we will revisit this topic in [11, 12].

## REFERENCES

- [1] H. S. Wilf, *generatingfunctionology*, 2<sup>nd</sup> ed., Academic Press, 1994, pp. 61–62; available online at <http://www.math.upenn.edu/~wilf/>; MR1277813 (95a:05002).
- [2] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A060594 and A114643.
- [3] Sloane, op. cit., A060839 and A160498.
- [4] S. Finch and P. Sebah, Squares and cubes modulo  $n$ , <http://arxiv.org/abs/math.NT/0604465>.

- [5] C. David, J. Fearnley and H. Kisilevsky, On the vanishing of twisted  $L$ -functions of elliptic curves, *Experiment. Math.* 13 (2004) 185–198; MR2068892 (2005e:11082).
- [6] Sloane, op. cit., A073103 and A160499.
- [7] S. R. Finch, Landau-Ramanujan constant, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 98–104; MR2003519 (2004i:00001).
- [8] D. Shanks and L. P. Schmid, Variations on a theorem of Landau. I, *Math. Comp.* 20 (1966) 551–569; MR0210678 (35 #1564).
- [9] P. Moree and H. J. J. te Riele, The hexagonal versus the square lattice, *Math. Comp.* 73 (2004) 451–473; preprint available online at <http://arxiv.org/abs/math.NT/0204332>; MR2034132 (2005b:11155).
- [10] G. H. Hardy, Ramanujan: *Twelve Lectures on Subjects Suggested by his Life and Work*, Chelsea, 1959, pp. 60–63; MR0106147 (21 #4881).
- [11] S. Finch and P. Sebah, Residue of a mod 5 Euler product, <http://arxiv.org/abs/0912.3677>.
- [12] S. Finch, G. Martin and P. Sebah, Roots of unity and nullity modulo  $n$ , *Proc. Amer. Math. Soc.*, to appear; <http://www.math.ubc.ca/~gerg/papers/downloads/RUNM.pdf>.

Steven Finch  
Dept. of Statistics  
Harvard University  
Cambridge, MA, USA  
*Steven.Finch@inria.fr*